

Superluminal Tunneling Confronts Special Relativity

Günter Nimtz

II. Physikalisches Institut, Universität zu Köln
Zülpicherstrasse 77, 50936 Köln (g.nimtz@uni-koeln.de)

Abstract

Experiments with evanescent modes and tunneling particles have revealed several non classical properties. The most important ones will be introduced:

- i) their signal velocity may be faster than light (i.e. superluminal),
- ii) they are described by virtual particles,
- iii) they are nonlocal and act at a distance,
- iv) the tunneling time is due to scattering time at the barrier front, there is zero time spent inside the barrier,
- v) this strange behavior is a universal field independent property. Actually, experimental tunneling data of phonons, photons, and electrons display a universal scattering time at the tunneling barrier front. The scattering time equals approximately the wave packet's oscillation time. Scattering time data from 10^{-3} s to 10^{-18} s have been measured and predicted.
- vi) the properties of evanescent, i.e. of tunneling modes are not compatible with the special theory of relativity.

References

- [1] T. Hartman, J. App. Phys., **33**, 3427-3433 (1962)
- [2] W. Franz, phys. stat. sol. **22**, K139 (1967)
- [3] C. Carniglia and L. Mandel, Phys. Rev. D, **3**, 280-296 (1971)
- [4] S. T. Ali, Phys.Rev.D, **7**, 1668-1675(1973)
- [5] F. Low and P. Mende, Annals of Physics (NewYork), **210**, 380-387 (1991)
- [6] A. Enders and G. Nimtz, J. Phys. I, (France) **2**, 1693-1698 (1992)
- [7] S. Esposito, Phys. Rev. E, **64**, 026609-1-8 (2001)
- [8] A. Haibel, G. Nimtz, Ann. Phys. (Leipzig) **10**, 707-712 (2001)
- [9] S. Longhi et al., Phys. Rev. E **64**, 055602-4 (2001)
- [10] G. Nimtz, Lecture Notes in Physics, **702**, 506-531 (2006)
- [11] A. A. Stahlhofen and G. Nimtz, Europhys. Letters, **76**, 189-195 (2006).
- [12] P. Eckle et al., Science **322**, 1525-1529 (2008)
- [13] G. Nimtz, Found. Phys. **39**, 1346-1355 (2009)
- [14] G. Nimtz, Found. Phys. **41**, 1193-1199 (2011)